

Home Search Collections Journals About Contact us My IOPscience

The Haber Bosch–harmful algal bloom (HB–HAB) link

This content has been downloaded from IOPscience. Please scroll down to see the full text.

View the table of contents for this issue, or go to the journal homepage for more

Download details:

IP Address: 50.138.226.125 This content was downloaded on 29/03/2015 at 22:03

Please note that terms and conditions apply.

Environ. Res. Lett. 9 (2014) 105001 (13pp)

The Haber Bosch–harmful algal bloom (HB–HAB) link

Patricia M Glibert¹, Roxane Maranger², Daniel J Sobota³ and Lex Bouwman^{4,5}

¹University of Maryland Center for Environmental Science, Horn Point Laboratory, PO Box 775, Cambridge, MD 21613, USA

² Groupe de Recherche Interuniversitaire en Limnologie et en Environnement Aquatique (GRIL), Département de Sciences Biologiques, Université de Montréal, Case Postale 6128, Succ. Centre-Ville, Montréal, Québec H3C 3J7, Canada

³Environmental Solutions Division, Oregon Department of Environmental Quality, 811 SW 6th Ave., Portland, OR 97206, USA

⁴ Department of Earth Sciences—Geochemistry, Faculty of Geosciences, Utrecht University, PO Box 80021, 3508 TA Utrecht, The Netherlands

⁵ PBL Netherlands Environmental Assessment Agency, PO Box 303, 3720 AH Bilthoven, The Netherlands

E-mail: glibert@umces.edu

Received 4 August 2014, revised 24 August 2014 Accepted for publication 28 August 2014 Published 6 October 2014

Abstract

Large-scale commercialization of the Haber-Bosch (HB) process is resulting in intensification of nitrogen (N) fertilizer use worldwide. Globally N fertilizer use is far outpacing that of phosphorus (P) fertilizer. Much of the increase in N fertilizers is also now in the form of urea, a reduced form of N. Incorporation of these fertilizers into agricultural products is inefficient leading to significant environmental pollution and aquatic eutrophication. Of particular concern is the increased occurrence of harmful algal blooms (HABs) in waters receiving nutrient enriched runoff. Many phytoplankton causing HABs have physiological adaptive strategies that make them favored under conditions of elevated N: P conditions and supply of chemically reduced N (ammonium, urea). We propose that the HB-HAB link is a function of (1) the inefficiency of incorporation of N fertilizers in the food supply chain, the leakiness of the N cycle from crop to table, and the fate of lost N relative to P to the environment; and (2) adaptive physiology of many HABs to thrive in environments in which there is excess N relative to classic nutrient stoichiometric proportions and where chemically reduced forms of N dominate. The rate of HAB expansion is particularly pronounced in China where N fertilizer use has escalated very rapidly, where soil retention is declining, and where blooms have had large economic and ecological impacts. There, in addition to increased use of urea and high N: P based fertilizers overall, escalating aquaculture production adds to the availability of reduced forms of N, as does atmospheric deposition of ammonia. HABs in both freshwaters and marginal seas in China are highly related to these overall changing N loads and ratios. Without more aggressive N control the future outlook in terms of HABs is likely to include more events, more often, and they may also be more toxic.

S Online supplementary data available from stacks.iop.org/ERL/9/105001/mmedia

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Keywords: nitrogen, eutrophication, nitrogen fertilizer, N:P ratio, cyanobacteria, dinoflagellates, harmful algal blooms

The industrial fixation of nitrogen gas (N_2) to ammonia (NH_3) , the Haber–Bosch (HB) process

$$N_2 + 3 H_2 - > 2NH_3$$

is considered to be one of the most important chemical reactions in the world (e.g., Smil 2001) and 'the greatest single experiment in global geo-engineering ever made' (Sutton *et al* 2013, p. 4). This reaction has produced the nitrogen (N) fertilizers that have contributed to the 'green revolution' responsible for increased food production that has supported the expansion of human population from ~2 billion in the early 20th century to >7 billion people today (Smil 1999, Erisman *et al* 2008).

Prior to World War II, the creation of reactive N was largely due to natural processes, including biological N fixation and lightning, and population expansion kept pace with its creation (Galloway *et al* 2002). After the mid-1940s and the commercialization and scaling up of the HB process, the manufacture and use of N expanded rapidly, from <10 MT N yr⁻¹ in 1950 to >170 MT N yr⁻¹ in 2013 (MT = megatonnes; figure 1(A); Constant and Sheldrick 1992, FAO 2012, Heffer and Prud'homme 2013). In fact, 85% of all synthetic N fertilizers have been created since 1985 (Howarth 2008).

The rate of change in use of N fertilizers has eclipsed that of phosphorus (P) fertilizers in large part due to this largescale capacity for anthropogenic synthesis. Global use of N fertilizer has increased nine-fold, while that of P has increased three-fold (Sutton *et al* 2013; figure 1(B)) and the upward trends in N: P in fertilizer application (figure 1(C)) are apparent in virtually all regions of the globe (figures 1(D)–(F)). In the US, it has been estimated that there has been at least a five-fold increase in reactive N use on average compared to pre-industrial time (Houlton *et al* 2013), but this increase is spatially variable ranging from negligible to 35-fold in different areas (Sobota *et al* 2013).

Although the HB process is the conversion of N2 to NH3, nearly 60% of all N fertilizer now used throughout most of the world is in the form of urea (CO(NH₂)₂; Constant and Sheldrick 1992, Glibert et al 2006, IFA 2014; figure 1(A). Urea production is an extension of the HB process, being produced by reacting CO₂ with anhydrous NH₃ under pressure at high temperatures. World use of urea as a fertilizer and feed additive has increased more than 100-fold in the past four decades (Glibert et al 2006). From 2001 to 2010, global urea use grew on average at a rate of $3.8\% \text{ yr}^{-1}$, and it is projected that from 2012 to 2017 an estimated 55 new urea manufacturing plants will be constructed worldwide, with half of these located in China (Heffer and Prud'homme 2013), contributing to another anticipated doubling by 2050 (Glibert et al 2006). Multiple factors, including the less explosive nature of urea relative to ammonium-nitrate fertilizer (NH₄NO₃), make the transportation and storage of this synthetically produced N form much safer and the preferred choice for agricultural applications.

The incorporation of agricultural N into plant biomass is extremely inefficient. Although the efficiency of N use in experimental fields may be much higher than the global average of 50% (Balasubramanian et al 2004), under practical conditions it is difficult to equate the N supply from fertilizer and from soil organic matter mineralization with the dynamics of crop N uptake demand (Dobermann and Cassman 2005). Considering the complete food chain, only ~10-30% of N applied actually reaches human consumers (Galloway et al 2002, Houlton et al 2013). The difficulty in improving N use efficiency in agriculture lies in the high mobility of N in the soil-plant system, and the variety of potential loss pathways ranging from NH₃ volatilization, denitrification, leaching and runoff, and other N transformation processes (Bouwman et al 2009). Urea inputs are typically hydrolyzed to NH⁺₄ in soil, but losses via volatilization and from runoff can be large and depend on the timing of application, weather, soil temperature and pH and other factors (Khakural and Alva 1995, Wali et al 2003). In regions such as China where the rate of fertilization has risen rapidly, the rate of soil retention of the excess N is actually declining (Cui et al 2013), leading to further environmental leakage.

The recovery of fertilizer P in crop products is also low (Syers *et al* 2008) but its biogeochemistry leads to proportionately greater retention within soils and sediments than N (e.g., Rhue and Harris 1999, Smil 2000, Bouwman *et al* 2009). The accumulation of residual soil P due to large fertilizer P surpluses over crop uptake during the 1960s, 1970s, and 1980s has led to an increased pool of plant available P in soils of most industrialized countries; a similar development was seen in later decades in India and China (Sattari *et al* 2012, 2014).

Among the various fates of the 'leaked N' are pathways that ultimately lead to N enrichment of lakes, rivers, and coastal waters. The major pathways for this leaked N include direct runoff, estimated to range up to 40% of inputs in large rivers (e.g., Howarth et al 2006) and atmospheric volatilization of NH⁺₄, and together these pathways can comprise more than half of the N input (e.g., Galloway et al 2004). P also runs off to receiving waters, but given the aforementioned change in patterns of fertilizer use and its biogeochemistry, the stoichiometry of the runoff has also changed in the last decades, leading to increasing N: P in receiving waters (e.g., Glibert et al 2013). It has been estimated that the atmospheric deposition of nutrients in the ocean is now ~ 20 times the Redfield ratio for N:P (Jickells 2006, Peñuelas et al 2012) and these changes are also having large consequences for N: P stoichiometry in lakes (Elser et al 2009). This change in stoichiometry has been further compounded since the mid-1980s and 1990s when the major industrialized nations began curtailing P use by removing it from detergents and by upgrading sewage treatment processes which generally are



Year

Figure 1. N and P (as P_2O_5) fertilizer use and change in N : P ratio of fertilizer use by weight for the world ((A), (B), and (C) respectively) and for selected countries or regions ((B), (D), and (E) respectively). Superimposed on the world N use graph (A) is the fraction of N use as urea (bars). Total N and P data are from FAO (2012) and data are the average of the three preceding years for each 5 yr period; urea data are from Constant and Sheldrick (1992) through 1990 and estimated at 3.8% growth per year thereafter, comparable to urea data reported from IFA (2014).

more efficient in removing P than N (Litke 1999, Van Drecht *et al* 2009).

In general, urea concentrations in aquatic ecosystems are less than those of the inorganic N forms NO_3^- and NH_4^+ , but depending on proximity of the land source to the water body, concentrations of urea in lacustrine, estuarine and coastal waters may be high, particularly when runoff occurs from heavily fertilized areas (Glibert et al 2005, 2006). Concentrations up to 25-50 µMN have been reported in tributaries of the Chesapeake Bay (Lomas et al 2002, Glibert et al 2005), and nearshore waters adjacent to the heavily fertilized Yaqui Valley, Mexico (Glibert et al 2006), among other coastal areas (Kudela et al 2008, Switzer 2008). Urea concentrations vary from undetectable to $150 \,\mu\text{M}\,\text{N}$ in Lake Kinneret, Israel (Berman 1974), in Polish lakes (Siuda and Chrost 2006) and in lakes of Central Canada where these concentrations also represented 10-50% of bioavailable N (Bogard et al 2012). Urea is also part of the dissolved organic N pool that is typically a very large component of the available N in nutrient rich waters (Glibert et al 2006, Solomon et al 2010).

Eutrophication is the process by which waters are enriched in nutrients, leading to effects such as increased algal growth and development of high biomass blooms, changes in species diversity of both the primary and secondary producers, reductions in dissolved oxygen, fish kills, and the increased frequency of harmful algal blooms (HABs; Nixon 1995, Cloern 2001). HABs are those proliferations of algae that can cause ecological harm to food webs when they accumulate in massive quantities, and they can cause ecological, human health, and economic impacts when these cells produce toxic or other bioactive compounds and when the decay of high algal biomass results in hypoxia (Hallegraeff 1993, Glibert et al 2005, Backer and McGillicuddy 2006). The most common HABs are either dinoflagellates or cyanobacteria, although not all dinoflagellates or cyanobacteria are harmful, and not all HABs are made up of these species groups. Cyanobacteria are the HAB functional group of proportionately greater concern in freshwater, while dinoflagellates are the HAB functional group of greater concern in estuarine and marine waters. HABs have been expanding globally, in spatial extent, in duration of blooms, and in intensity (Anderson et al 2002, 2008, Glibert et al 2005) and a critical question has been the extent to which this change is associated with eutrophication and/or accelerated by climate or other factors (e.g., Paerl and Scott 2010). There are many reports of increases in HABs associated with eutrophication or nutrient loading (e.g., Anderson et al 2002, 2008, Glibert et al 2006, 2010, Heisler et al 2008), but the complexity of the relationship is far from understood. Many nutrient reduction strategies are focused on reducing algal biomass (e.g., total chlorophyll), while appropriate efforts or best management practices that may specifically prevent toxic HABs have remained much more challenging (Jewett et al 2008).

One of the most central tenets of aquatic science is that algal biomass and production in lakes and other freshwaters is limited by the availability of P, while that in marine waters is more often limited by the availability of N (e.g., Ryther and Dunstan 1971, Schindler 1977). However, P limitation in lakes is not universal (Lewis and Wurtsbaugh 2008) as some regions are either naturally (Finlay et al 2010) or culturally enriched (Bennett et al 2001) in P relative to N, and excess N loading is changing the nutrient stoichiometry and limiting element in some coastal areas (e.g., Sylvan et al 2006). It is commonly assumed that to control eutrophication the only focus should be on that nutrient which is classically considered 'limiting.' Such an argument is typically extended to promote enhanced P control over N control (e.g., Schindler et al 2008, Wang and Wang 2009) for multiple reasons, among which it is often assumed that N2-fixing toxic cyanobacteria will be favored when N is limiting. Contrary to this common perception, in enriched systems, some toxic cyanobacteria are more common under elevated N: P conditions or when total N (TN) concentrations are high, and several cross comparative studies show that total cyanobacterial biomass can be predicted from increasing TN concentrations and from changes in TN to total P ratios (Smith 1983, Downing et al 2001, Kosten et al 2012, Dolman et al 2012). Therefore, species composition is important to consider (Dolman et al 2012) as, in fact, some of the most toxic cyanobacteria, namely Microcystis sp., are not N2-fixing species.

Further arguments for P relative to N control are that P does not have a gaseous form and therefore cannot be permanently removed from lakes, whereas N can be lost to the atmosphere through denitrification provided conditions are optimal to favor coupled nitrification–denitrification (Seitzinger 1988, Cornwell *et al* 1999). However, rates of N₂-fixation do not necessarily offset N limitation (Scott and McCarthy 2010, Lewis *et al* 2011) and not all N enrichment is removed through denitrification resulting in eutrophication related problems that may be spatially and temporally displaced from the nutrient source (Conley *et al* 2009, Pearl 2009, Glibert *et al* 2011).

Here we make the case that accelerated P reduction over N control is insufficient in reducing the frequency of HABs; we base this on our growing understanding of HAB physiology and we support this with long-term trend data from one of the most eutrophic regions of the world, China. We also argue that increasing N:P environments further favor HABs when the N form is disproportionately in chemically reduced form (i.e. urea, NH₄) relative to chemically oxidized form (i.e., NO_3^-). We thus propose that the HB-HAB link is a function of (1) the inefficiency of incorporation of N fertilizers in the food supply chain, the leakiness of the N cycle from crop to table, and the fate of the lost N to the environment as described above; and (2) adaptive physiology of many HABs to thrive in environments in which there is excess N relative to classic nutrient stoichiometric proportions and where chemically reduced forms of N are increasing. We review physiological evidence for this at the cellular level, and we highlight long-term trends in HABs and fertilizer in China demonstrating increased HAB proliferations for systems in which N:P stoichiometry has been altered due to continued N loading with P reduction and the greater proportional use of urea as the main N fertilizer.

There are a number of specific physiological strategies that allow certain types of algae to thrive under conditions of elevated N: P availability relative to classic Redfieldian proportions (Redfield 1934), but not all cells necessarily have all such adaptive strategies (Glibert and Burkholder 2011). The first strategy is a low overall requirement for P. Very small cells, such as picocyanobacteria, have a lower requirement for P due to the smaller need for structural components in the cell (Finkel et al 2010). The second strategy is the ability to 'make do with less' which may be accomplished by physiological substitution of a P-containing lipid with a non-P-containing lipid (sulfolipid), and many cyanobacteria are able to do this (Van Mooy et al 2009). Thus the cellular C: P content of Synechococcus is about 100, whereas this ratio in a typical diatom is about 50 (Finkel et al 2010). The third strategy is the ability to acquire P in organic or particulate form, via alkaline phosphatase activity or mixotrophy, which may provide some cells a source of P not available to those cells dependent on inorganic P for their nutrition. Many dinoflagellates have a comparatively high cellular P requirement, and therefore the ability to consume particulate P may be an important reason why these types of cells can thrive when some others cannot. An added competitive benefit for these cells is that there may also be a growth advantage when feeding mixotrophically, compared to pure autotrophic growth (Jeong et al 2004, Glibert et al 2009, Flynn et al 2013), thus mixotrophy is a major mode of nutrition by HABs in eutrophic waters (Burkholder et al 2008) and may help to sustain blooms when dissolved nutrients are depleted.

Of particular concern is the association of increased toxin in many HABs under condition of elevated N : P availability. Many cyanobacteria and dinoflagellate toxins are N-rich compounds and thus these cells require a supply of N in order to synthesize these metabolites. The most ubiquitous cyanotoxin in freshwater systems are microcystins (MCs), hepatotoxic compounds that can be lethal to mammals if ingested (Carmichael 1994, Chorus and Bartram 1999), some congeners of which are significantly more toxic than others (Sivonen and Jones 1999). Microcystis (Chroococcales), Anabaena (Nostocales), and Planktothrix (Oscillatoriales) are among the taxa that can synthesize MCs (Cronberg and Annadottir 2006). Total concentrations of MC have been strongly related to N concentrations in several comparative studies (Giani et al 2005, Rolland et al 2005, Dolman et al 2012, Monchamp et al 2014). Although negative relationships between MC and N: P ratios have been reported for lake Tai (Taihu), China (Otten et al 2012), but in a cross system analysis of freshwaters (Orihel et al 2012), the authors of the findings of negative correlations emphasize that such results are relevant only to hypereutrophic conditions in which TN concentrations are also high (>100 μ M N). Furthermore, a reanalysis of the aforementioned comparative study reported highest MC concentration at intermediate N : P ratios (Scott et al 2013). Excess N and high N: P ratios have also been related to MC production under controlled culture conditions (e.g., Lee et al 2000, Oh et al 2000, Vézie et al 2002, Downing et al 2005, Van de Waal et al 2009). Some in situ evidence, albeit weak, suggests that increased N availability may influence the MC congener type to more toxic variants that have higher N content (e.g., Van de Waal *et al* 2009) and that P limitation causes an increase in N-rich toxins of numerous HABs (Van de Waal *et al* 2014).

Some marine HABs also show increased toxin production under conditions of elevated N: P ratios. As examples, under conditions of elevated N: P, hemolytic activity per cell increases by up to an order of magnitude in the prymnesiophytes *Prymensium parvum* and *Chrysochromulina* (now *Prymnesium*) polylepis (Johansson and Granéli 1999), and neurotoxin production increases in the diatom *Pseudo-nitzschia multiseries* and in the dinoflagellates *Karlodinium venificum*, *Alexandrium* sp., and *Karenia brevis* (Granéli and Flynn 2006, Hardison *et al* 2013).

Recent reviews of the physiological bases of N uptake as well as molecular and metatranscriptomic data lend considerable support to the emerging conclusion that diatoms are specialists in use of oxidized forms of N, while cyanobacteria and dinoflagellates are specialists in reduced forms of N (e.g., Glibert et al 2014a and references therein). A considerable amount of experimental evidence supports the notion that freshwater cyanobacteria seem to favor reduced N forms (Blomqvist et al 1994, Berman and Chava 1999). Extensive laboratory molecular evidence of the use of both urea and NH₄⁺ by cyanobacteria (Flores and Herrero 2005), and gene expression data (Ginn et al 2009) also lend support to this conclusion. Mesocosm studies conducted in fresh and brackish systems further illustrate that when enriched with oxidized versus reduced forms of N, even when the TN supply remains the same, proportionately more diatoms are produced under oxidized N conditons while more cyanobacteria and cryptophytes are produced under conditions of increasing reduced forms of N (e.g., Finlay et al 2010, Donald et al 2013, Glibert et al 2014a). Donald et al (2011) found in mesocosm enrichment studies conducted in the Northern Prairie lakes of Canada, that urea additions stimulated M. aeruginosa growth but MC concentrations increased even more. Of the comparative field studies in freshwater, toxic cyanobacterial species appear to be favored over diatoms as a function of N availability in reduced relative to more oxidized forms in the hypereutrophic Taihu and Lake Okechobee (McCarthy et al 2009). In marine systems, cyanobacteria and dinoflagellates have also been associated with proportionately greater use of reduced forms of N and diatoms greater use of oxidized forms of N (e.g., Berg et al 2003, Heil et al 2007).

In situ evidence of the role of N forms in cyanobacterial community composition remain rare and influence on MC concentrations or congener composition even rarer. In Québec, Canada, the ratio of N : P in fertilizer acquisition increased steadily from 1977 to 1995, and accelerated from 1995 to peaks of >3 (on a weight basis) from 2006 to 2008 (figure 2(A)). There was a marked increase in the purchase of N fertilizer in the form of urea or products containing urea as of 2003 with a urea peak observed in 2007. That year was considered an 'exceptionally favorable' year for agricultural commodities (Heffer and Prud'homme 2008), and a sustained higher overall use of urea products has occurred since that time (figure 2(B)). Records of reported cyanobacterial incidences in lakes across



Figure 2. (A) N : P ratio of fertilizer use (by weight) in Québec since 1967 illustrating year-to-year variability superimposed on long-term trends. (B) Trends in use of different N fertilizer types since 1998 in Québec (urea-squares; NH_4NO_3 -circles; urea $-NH_4-NO_3$ (UAN)-triangles); note the increasing use of UAN versus the more consistent use of urea and NH_4NO_3 . (C) Number of lakes with reported cyanobacteria blooms from over 450 systems surveyed in Québec since 2004. Note the time scale zooms in to progressively recent years from (A) to (C); also note the peaks in all panels in the year 2007. (D) Correlation between the number of lakes reporting cyanobacterial blooms from 2004 to 2012 and the use of urea plus combined N (UAN).

the province (see SI) suggest that the peak in events occurred in 2007 (figure 2(C)), when both the N : P ratio and urea use were high. Since then the incidences of blooms have remained high (figure 2(C)). The correlation of incidences of blooms in these lakes with the increase in forms of N fertilizer containing urea (figure 2(D); r^2 =0.42, p=0.058) is suggestive of such an effect, although more years of data will be required to substantiate this relationship. While these patterns provide only indirect evidence, in a comparison over the course of the growing season of three lakes in Québec known to have toxic cyanobacteria, cyanobacterial community structure was

primarily influenced by the availability of chemically reduced and organic N forms (DON and NH_4^+) and temperature (Monchamp *et al* 2014). N forms and concentration however did not influence congener composition or the toxicity of the dominant variant in that study, but cyanobacterial community structure did. This suggests complex interaction between the availability of N and other environmental variables in influencing community structure with an indirect effect on congener composition and overall bloom toxicity.

Several of the more potent marine HAB species show the same trend. For Alexandrium tamarense, the availability of urea has also been related to toxin content of the cells: the toxin content for urea-grown cells was found to be higher than that of NO₃-grown cells, but not as high as cells grown on NH_4^+ (Leong *et al* 2004). Furthermore the biosynthesis of toxin when grown on urea appears to differ from that which occurs under NO₃⁻ or NH₄⁺ growth conditions. For another dinoflagellate, Karenia brevis, up to six-fold increases in toxin content have been observed during growth with elevated urea availability compared to controls without urea enrichment (Shimizu et al 1993). For the toxic diatom, Pseudo-nitzschia sp., increases in toxicity in both laboratory cultures and natural field assemblages have also been found for cells growing on urea compared to those growing on NH₄⁺ or NO₃ (Cochlan et al 2008, Kudela et al 2008). On a global scale there is a simultaneous increase of total fertilizer N, particularly in the form of urea, and the frequency and extent of a number of HAB cyanobacteria and dinoflagellate species providing further, though indirect, evidence for a relationship between urea and HABs (Glibert et al 2006, 2008).

China presents an interesting case study in terms of the relationships between increasing HAB frequency and the changes in fertilizer use and export. All data used herein come from pre-existing data sources, industry statistics and published literature (see SI). Fertilizer N use in China has escalated from about ~0.5 MT in the early 1960s to 42 MT around 2010, with the fraction of urea increasing nearly five-fold over just the past two decades (figure 3(A); IFA 2014, FAO 2012, Zhang and Zhang, no date). River export of N increased from 1980 to 2010 from \sim 500 to >1200 kg N km⁻² yr⁻¹ in the Changjiang River (Yantgtze River), from ~100 to \sim 200 kg N km⁻² yr⁻¹ in the Huanghe River (Yellow River), and from ~ 400 to > 1200 kg N km⁻² yr⁻¹ in the Zhujiang River (Pearl River) basins, the latter having one of the largest N: P ratios in the world (Ti and Yan 2013). The annual N load from the Changjiang River to the coastal ocean is higher than loads from the Mississippi and the Amazon rivers (Goolsby and Battaglin 2001, Duan et al 2008). Atmospheric sources of NH₄⁺, which is the dominant form of inorganic N on a regional basis, and which exceeds that of oxidized N forms in the Changjiang River basin, are due primarily to livestock excretion, fertilizer N use, and human waste (52%, 33% and 13% respectively, Xiao et al 2010).

The number of HABs has increased in all waters of China in the past three decades and most inland and coastal waters are rated in the moderately to severely polluted range (Wang *et al* 2011, Ti and Yan 2013). In Taihu, the largest lake in the Changjiang watershed, the duration of cyanobacterial



Figure 3. (A) Comparison of N (red circles) and P (blue squares) fertilizer use and the use of urea (black crosses) in China from 1975 to 2005. N and P data are from FAO (2012) while urea data are from Zhang and Zhang (no date) and represent the total of production and imports minus exports. Also shown (gray triangles) is the calculated reactive N retention capacity of the Chinese landscape as estimated by Cui *et al* (2013). (B) Change in annual duration of *Microcystis* blooms in Lake Tai (Taihu) in months, urea fertilizer use scaled to that in the Changjiang watershed and the ratio of use of urea : P_2O_5 fertilizer. *Microcystis* data are from Duan *et al* (2009) reprinted with permission of the American Chemical Society, and total urea and P_2O_5 data for China are from Zhang and Zhang (no date). (C) and (D) Correlations between annual duration of blooms (months) and urea use in Changjiang watershed (C) and the urea : P_2O_5 ratio (D).

Microcystis blooms has increased from ~1 month yr⁻¹ to nearly 10 months yr⁻¹ in the past 15 yr (Duan *et al* 2009; figure 3(B)). The Changjiang watershed produces >30% of China's agricultural output (Xing and Zu 2002) and in the Taihu region, N fertilizers from manures and mineral fertilizers have increased eight- to ten-fold since the 1950s (Chen *et al* 2008) and fertilizer N use in Jiangsu Province (2.5 MT N yr⁻¹) is 80% of TN inputs (PBL 2012). The change in HABs in Taihu is strongly related to the increase in urea and in the ratio of urea : P₂O₅ use scaled to the Changjiang watershed (Chen *et al* 2003, Ye *et al* 2007, Duan *et al* 2009; figures 3(B)–(D); r^2 =0.85 and 0.92, respectively, p<0.01).

These trends in availability of reduced forms of N and in increased N : P ratios in this region are further accelerated by aquaculture development and atmospheric deposition. In 2007, 13% of Chinese inland aquaculture production was in Jiangsu (China Fishery Yearbook 2007), and large quantities of NH_4^+ and urea with high N : P ratio are also discharged from that source to Taihu. It should be noted that waste from aquaculture not only has a high N : P ratio (>20) but is largely in reduced N form with significant amounts of urea (Bouwman *et al* 2013). Also, 80% of the atmospheric deposition in this region has been associated with NH_4^+ mobilization from intense fertilization during the growing period (Chen



Figure 4. (A) The total number of recorded HAB occurrences (red squares) along the coast of China and their aerial extent (open triangles). Data are from Wang *et al* (2011), originally reported by SOA (2010; reproduced under Creative Commons license). (B) and C Correlations between annual number of HABs or areal extent of HABs in China marginal seas and urea use. (D) The annual number of HAB occurrences in the Huanghai (Yellow) Sea region (red squares) and the molar N : P ratio of southern Huanghai Sea in spring (blue diamonds). Data are from Fu *et al* (2012), with the HAB data originally reported by SOA (data reproduced with the permission of the Chinese Society of Oceanography and Springer-Verlag). (E) The annual number of HAB occurrences in the Northern South China Sea (red squares) and the molar N : P ratio in the water column. Data on HABs are from Wang *et al* (2008) and N : P data are from Ning *et al* (2009; reproduced under creative commons license) and represent the average values for the upper 200 m of the water column.

et al 2008). With intensive agriculture, N deposition onto the lake is a direct N input not accompanied by P.

In Chinese marginal seas there are similar trends of increased HABs over the past several decades (Wang *et al* 2008, 2011; figure 4(A)). When all China's marginal sea bloom occurrences are considered, there are strong relationships between both number of HAB reports and aerial HAB extent and N use as urea (figures 4(B) and (C); $r^2 = 0.53$ and 0.37, respectively, p < 0.01).

In both the Huanghai (Yellow) Sea region and in the Northern South China Sea, the number of reported occurrences of HABs has increased in parallel with the long-term trend in N:P of the water column (figures 4(D) and (E); $r^2 = 0.46$ and 0.29, respectively, p = 0.06). As in the Taihu example, these relationships do not account for nutrient input from coastal aquaculture or from atmospheric NH₄⁺ deposition that would enhance these relationships. Coastal N release from Chinese mariculture increased from 2000 to 2010 by 45% on average and up to 63% in some provinces while P increase was only 37% (Bouwman *et al* 2013). In 2004, NH₄⁺ and organic N comprised 75% of N in wet atmospheric deposition in Southeast China (Chen *et al* 2011).

At the mouth of the Changjiang River, molar $NO_3^-: PO_4^{-1}$ values were ~30–40 in the 1960s but had risen to >250 by the late 1990s (Shen and Liu 2009; figure 4(D)). Occurrences of HABs in the East China Sea were rare in the 1970s, had increased to a few dozen over the 1980s but had increased to >130 in just the 5 yr from 1992 to 1997 (Shen and Liu 2009) but the scale of the blooms soared in the years since; the spatial scale of the annual blooms increased from 1000s of km² in 2000 to >15 000 km² by 2005 with many millions of dollars lost in high value aquaculture products due to associated fish kills (Li *et al* 2009). The dinoflagellates *Prorocentrum donghaiense* and *Karenia mikimotoi* are among the common HABs now reported in East China Sea (e.g., Zhou *et al* 2008, Li *et al* 2009).

In the Huanghai Sea region, inorganic N : P ratios are now about twice Redfield proportions, and about four-fold higher than in the 1990s (figure 4(E)). There has also been nearly a six-fold increase in HAB occurrences and a shift to proportionately more dinoflagellates compared to diatoms (Fu *et al* 2012). In the South China Sea region, water column inorganic N : P ratios increased from ~2 in the mid-1980s to >20 in the early 2000s (Ning *et al* 2009). In addition to the increase in number of HABs, a change in species composition to increasing dominance of species such as *Chattonella*, *Gymnodinium breve*, and *Dinophysis* has occurred (Wang *et al* 2008).

The results herein lend support the view that both N and P controls are necessary to reduce eutrophication in both freshwater and marine waters (e.g., Burkholder *et al* 2006, Howarth and Paerl 2008, Conley *et al* 2009, Paerl 2009, Glibert *et al* 2011, 2013). Loss of biodiversity, and effects on ecosystem and human health due to eutrophication are considered major challenges of our current day (e.g., Borja 2014). Given the known inefficiencies in all aspects of N use in both industrialized nations and throughout the world, the benefits of increased emphasis on N reduction, and improved N use

efficiencies at all levels of the production side, would have far reaching benefits to ecosystems and especially to water quality (Houlton et al 2013, Sobota et al 2013). Calculated damage costs of loss of ecosystem services, eutrophication and human health due to loss of reactive N to the environment are large (Compton et al 2011, Sobota et al this issue). So too are the economic impacts to aquaculture, increased risks to human health and ecosystems, and losses to fisheries and ecosystem services due to HABs (Hoagland and Scatasta 2006). More work is needed to accurately quantify the sources and fluxes of N at all stages of the supply-to-loss pathways, and there is no question that control of N may be more challenging than control of P. Given our increasing knowledge of the physiological response of many HABs to increasing N: P and to increasing ratios of reduced : oxidized forms of N, we can conclude that without more aggressive N control the future outlook in terms of HABs is likely to include more events, more often, and such events may also be more toxic (O'Neil et al 2012, Glibert et al 2013, 2014b, Paerl and Otten 2013). Indeed, a recent modeling effort in which the physiological responses of HABs to altered N:P ratios and altered N form, even without any further increases in N loading, in conjunction with projected climate change effects, suggest an expansion in area and/or number of months annually conducive to development of several HAB genera along the NW European Shelf-Baltic Sea system and NE Asia by end of the century (Glibert et al 2014b). Such projections alone should be cause for advancing our understanding of the relationships between HABs and nutrient loading, and together with the projected global expansion in N loading should be serve to sound the alarm that our existing approaches to nutrient management of sensitive coastal and freshwaters are not sufficient, particularly in the face of climate change and other stressors.

Acknowledgements

Discussions for this paper were initiated at the 6th International Nitrogen Conference in Kampala, Uganda, 17–23 November 2013 based on presentations by PMG, RM, and DJS. This is a contribution of the GEOHAB core research project on HABs and eutrophication, and of SCOR Working Group 132 on Land Based Nutrient Pollution and Harmful Algae, funded through the Scientific Committee on Oceanic Research and the Land Ocean Interaction at the Coastal Zone Program, and of the Groupe de recherche interuniversitaire en limnologie et environment aquatique (GRIL). This is contribution number 4922 from the University of Maryland Center for Environmental Science.

References

Anderson D A, Burkholder J M, Cochlan W, Glibert P M, Gobler C, Heil C, Kudela R, Parsons T, Trainer V and Vargo G 2008 Harmful algal blooms in the United States: linkages to eutrophication *Harmful Algae* 8 39–53

- Anderson D, Glibert P M and Burkholder J M 2002 Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences *Estuaries* 25 704–26
- Backer L C and McGillicuddy D G Jr 2006 Harmful algal blooms at the interface between coastal oceanography and human health *Oceanography* **19** 94–106

Balasubramanian V, Alves B, Aulakh M, Bekunda M, Cai Z, Drinkwater L, Mugendi D, Van Kessel C and Oenema O 2004 Crop, environmental, and management factors affecting nitrogen use efficiency Agriculture and the Nitrogen Cycle ed A R Mosier, J K Syers and J R Freney (Washington, DC: Island Press) pp 19–33

Bennett E M, Carpenter S R and Caraco N F 2001 Human impact on erodible phosphorus and eutrophication: a global perspective *Bioscience* 51 227–34

Berg G M, Balode M, Purina I, Bekere S, Béchemin C and Maestrini S 2003 Plankton community composition in relation to availability an uptake of oxidized and reduced nitrogen *Aquat. Microb. Ecol.* **30** 263–74

Berman T 1974 Urea in waters of lake Kinneret (Sea of Galilee) Limnol. Oceanogr. **19** 977–80

Berman T and Chava S 1999 Algal growth on organic compounds as nitrogen sources J. Plankt. Res. 21 1423–37

Blomqvist P, Pettersson A and Hyenstrand P 1994 Ammonium–nitrogen—a key regulatory factor causing dominance of non-nitrogen-fixing cyanobacteria in aquatic systems Arch. Für Hydrob. 132 141–64

Bogard M J, Donald D B, Finlay K and Leavitt P R 2012 Distribution and regulation of urea in lakes of Central North America *Freshw. Biol.* **57** 1277–92

Borja A 2014 Grand challenges in marine ecosystems ecology Frontiers Mar. Sci. 1: article 1 doi:10.3389/fmars.2014.00001

Bouwman A F, Beusen A H W and Billen G 2009 Human alteration of the global nitrogen and phosphorus soil balances for the period 1970–2050 Glob. Biogeochem. Cycles 23 GBA04

Bouwman L, Beusen A, Glibert P M, Overbeek C, Pawlowski M, Herrera J, Mulsow S, Yu R and Zhou M 2013 Mariculture: significant and expanding cause of coastal nutrient enrichment *Environ. Res. Lett.* **8** 044026

Burkholder J M et al 2006 Comprehensive trend analysis of nutrients and related variables in a large eutrophic estuary: a decadal study of anthropogenic and climatic influences *Limnol. Oceanogr.* 51 463–87

Burkholder J M, Glibert P M and Skelton H 2008 Mixotrophy, a major mode of nutrition for harmful algal species in eutrophic waters *Harmful Algae* 8 77–93

Carmichael W W 1994 Toxins of cyanobacteria Sci. Am. 270 78-86

Chen N, Hong H, Huang Q and Wu J 2011 Atmospheric nitrogen depositon and its long-term dynamcis in a Southeast China coastal area *J. Environ. Manage.* **92** 1663–7

Chen N, Hong H, Zhang L and Cao W 2008 Nitrogen sources and exports in an agricultural watershed in Southeast China *Biogeochemistry* 87 169–79

Chen Y W, Fan C, Teubner K and Dokulil M 2003 Changes in nutrients and phytoplankton chlorophyll a in a large shallow lake, Taihu, China: an 8-year investigation *Hydrobiologia* **506–509** 273–9

China Fishery Yearbook 2007 China Ministry of Agriculture Bureau of Fisheries (Beijing: Chinese Fishery Yearbook Publishing House)

Chorus I and Bartram J 1999 *Toxic Cyanobacteria in Water: A Guide to their Public Health Consequences, Monitoring and Management* (London: Spon) p 416

Cloern J E 2001 Our evolving conceptual model of the coastal eutrophication problem *Mar. Ecol. Prog. Ser.* **210** 223–53

Cochlan W P, Herndon J and Kudela R M 2008 Inorganic and organic nitrogen uptake by the toxigenic diatom *Pseudonitzschia australis* (Bacillariophyceae) *Harmful Algae* 8 111–8 Compton J E, Harrison J A, Dennis R L, Greaver T L, Hill B H, Jordan S J, Walker H and Campbell H V 2011 Ecosystem services altered by human changes in the nitrogen cycle: a new perspective for US decision making *Ecol. Lett.* 14 804–15

Conley D J, Paerl H W, Howarth R W, Boesch D F, Seitzinger S P, Havens K E, Lancelot C and Likens G E 2009 Controlling eutrophication: nitrogen and phosphorus *Science* **323** 1014–5

Constant K M and Sheldrick W F 1992 *World Nitrogen Survey* Technical paper 174 (Washington, DC: World Bank)

Cronberg G and Annadottir H 2006 Manual on Aquatic Cyanobacteria, a Photoguide and A Synopsis of their Toxicity (Copenhagen: ISSHA) p 106

Cornwell J C, Kemp W M and Kana T M 1999 Denitrification in coastal ecosystems: environmental controls and aspects of spatial and temporal scale Aquat. Ecol. 33 41–54

Cui S, Shi Y, Grffman P M, Schlesinger W H and Zhu Y-G 2013 Centennial-scale analysis of the creation and fate of reactive nitrogen in China (1910–2010) *Proc. Natl Acad. Sci. USA* 110 2052–7

Dobermann A and Cassman K G 2005 Cereal area and nitrogen use efficiency are drivers of future nitrogen fertilizer consumption *Sci. China Ser.* C. *Life Sci.* **48** Supp 1–14

Dolman A M, Rücker J, Pick F R, Fastner J, Rohrlack T, Mischke U and Wiedner C 2012 Cyanobacteria and cyanotoxins: the influence of nitrogen versus phosphorus *PLoS One* 7 e38757

Donald D B, Bogard M J, Finlay K, Bunting L and Leavitt P R 2013 Phytoplankton-specific response to enrichment of phosphorusrich surface waters with ammonium, nitrate, and urea *PLoS One* 8 e53277

Donald D B, Bogard M J, Finlay K and Leavitt P R 2011 Comparative effects of urea, ammonium, and nitrate on phytoplankton abundance, community composition, and toxicity in hypereutrophic freshwaters *Limnol. Oceanogr.* 56 2161–75

Downing J A, Watson S B and Mccaule E 2001 Predicting cyanobacteria dominance in lakes *Can. J. Fish. Aquat. Sci.* 58 1905–8

Downing T G, Sember C S, Gehringer M M and Leukes W 2005 Medium N : P ratios and specific growth rate comodulate microcystin and protein content in *Microcystis aeruginosa* PCC7806 and *M. aeruginosa* UV027 *Microb. Ecol.* **49** 468–73

Duan H R, Ma X, Xu F, Kong F, Zhang S, Kong W, Hao J and Shang L 2009 Two decade reconstruction of algal blooms in China's Lake Taihu *Environ. Sci. Technol.* 43 3524–8

Duan S W, Liang T, Zhang S, Wang L J, Zhang X M and Chen X B 2008 Seasonal changes in nitrogen and phosphorus transport in the lower Changjiang River before the construction of the three gorges dam *Estuarine Coast. Shelf Sci.* **79** 239–50

Elser J J, Andersen T, Baron J S, Bergström A K, Jansson K M, Nydick K R, Steger L and Hessen D O 2009 Shifts in lake N : P stoichiometry and nutrient limitation driven by atmospheric nitrogen deposition *Science* **326** 835–7

Erisman J W, Sutton M A, Galloway J N, Klimont Z and Winiwater W 2008 How a century of ammonia synthesis changed the world *Nat. Geosci.* **1** 636–9

FAO 2012 FAOSTAT Database Collections (Rome: FAO) (http:// faostat.fao.org)

Finkel Z V, Beardall J, Flynn K J, Quiqq A, Rees T A and Raven J A 2010 Phytoplankton in a changing world: cells size and elemental stoichiometry J. Plankt. Res. 32 119–37

Finlay K, Patoine A, Donald D B, Bogard M J and Leavitt P R 2010 Experimental evidence that pollution with urea can degrade water quality in phosphorus-rich lakes of the Northern great plains *Limnol. Oceanogr.* 55 1213–30

Flores E and Herrero A 2005 Nitrogen assimilation and nitrogen control in cyanobacteria *Biochem. Soc. Trans.* **33** 164–7

Flynn K J, Stoecker D K, Mitra A, Raven J A, Glibert P M, Hansen P J, Granéli E and Burkholder J M 2013 Misuse of the phytoplankton–zooplankton dichotomy: the need to assign organisms as mixotrophs within plankton functional types *J. Plankt. Res.* 35 3–11

- Fu M, Wang Z, Pu X, Xu Z and Zhu M 2012 Changes in nutrient concentrations and N : P : Si ratios and their possible impacts on the Huanghai Sea ecosystem Acta Oceanol. Sin. 31 101–12
- Galloway J N et al 2004 Nitrogen cycles: past, present, and future Biogeochemistry 70 153–226
- Galloway J N, Cowling E B, Seitzinger S P and Socolow R H 2002 Reactive nitrogen: too much of a good thing? *Ambio* **31** 60–3
- Giani A, Bird D, Prairie Y and Lawrence J 2005 Empirical study of cyanobacterial toxicity along a trophic gradient of lakes *Can. J. Fish. Aquat. Sci.* **2109** 2100–9
- Ginn H, Pearson L and Neilan B 2009 Hepatotoxin biosynthesis and regulation in cyanobacteria—the putative involvement of nitrogen and iron homeostasis mechanisms *Chiang Mai J. Sci.* 36 200–23
- Glibert P M, Allen J I, Artioli Y, Beusen A, Bouwman L, Harle J, Holmes R and Holt J 2014b Vulnerability of coastal ecosystems to changes in harmful algal bloom distribution in response to climate change: projections based on model analysis *Glob. Change Biol.* doi:10.1111/gcb.12662
- Glibert P M, Allen J I, Bouwman L, Brown C, Flynn K J, Lewitus A and Madden C 2010 Modeling of HABs and eutrophication: status, advances, challenges J. Mar. Syst. 83 262–75
- Glibert P M, Anderson D M, Gentien P, Granéli E and Sellner K G 2005 The global, complex phenomenon of harmful algal blooms *Oceanography* 18 136–47
- Glibert P M and Burkholder J M 2011 Eutrophication and HABs: strategies for nutrient uptake and growth outside the redfield comfort zone *Chin. J. Oceanol. Limnol.* **29** 724–38
- Glibert P M, Burkholder J M, Kana T M, Alexander J A, Schiller C and Skelton H 2009 Grazing by *Karenia brevis* on *Synechococcus* enhances their growth rate and may help to sustain blooms *Aquat. Microb. Ecol.* 55 17–30
- Glibert P M, Fullerton D, Burkholder J M, Cornwell J and Kana T M 2011 Ecological stoichiometry, biogeochemical cycling, invasive species and aquatic food webs: San Francisco Estuary and comparative systems *Rev. Fish. Sci.* **19** 358–417
- Glibert P M, Harrison J, Heil C and Seitzinger S 2006 Escalating worldwide use of urea—a global change contributing to coastal eutrophication *Biogeochemistry* 77 441–63
- Glibert P M, Kana T M and Brown K 2013 From limitation to excess: consequences of substrate excess and stoichiometry for phytoplankton physiology, trophodynamics and biogeochemistry, and implications for modeling J. Mar. Syst. 125 14–28
- Glibert P M, Mayorga E and Seitzinger S 2008 *Prorocentrum minimum* tracks anthropogenic nitrogen and phosphorus inputs on a global basis: application of spatially explicit nutrient export models *Harmful Algae* **8** 33–8
- Glibert P M, Trice T M, Michael B and Lane L 2005 Urea in the tributaries of the Chesapeake and coastal bays of Maryland *Water Air Soil Pollut.* 160 229–43
- Glibert P M, Wilkerson F, Dugdale R C, Parker A E, Alexander J A, Blaser S and Murasko S 2014a Microbial communities from San Francisco Bay Delta respond differently to oxidized and reduced nitrogen substrates—even under conditions that would otherwise suggest nitrogen sufficiency *Frontiers Mar. Sci.* 1: article 17 doi:10.3389/fmars.2014.00017
- Goolsby D A and Battaglin W A 2001 Long-term changes in concentrations and flux of nitrogen in the Mississippi River basin, USA *Hydrol. Proc.* 15 1209–26
- Granéli E and Flynn K 2006 Chemical and physical factors influencing toxin content *Ecology of Harmful Algae* ed E Granéli and J T Turner (Dordrecht: Springer) pp 229–41
- Hallegraeff G M 1993 A review of harmful algal blooms and their apparent global increase *Phycologia* **32** 79–99

- Hardison D R, Sunda W G, Shea D and Litaker R W 2013 Increased toxicity of *Karenia brevis* during phosphate limited growth: ecological and evolutionary implications *PLoS One* **8** e58545
- Heffer P and Prud'homme M 2008 Outlook for world fertilizer demand, supply, and supply/demand balance *Turk. J. Agric. For.* 32 159–64
- Heffer P and Prud'homme M 2013 *Fertilizer outlook 2013–2017* 81st IFA Annual Conf. (Chicago (USA), 20–22 May 2013) (www.fertilizer.org) Accessed June 22, 2014
- Heil C A, Revilla M, Glibert P M and Murasko S 2007 Nutrient quality drives phytoplankton community composition on the West Florida shelf *Limnol. Oceanogr.* 52 1067–78
- Heisler J et al 2008 Eutrophication and harmful algal blooms: a scientific consensus Harmful Algae 8 3-13
- Hoagland P and Scatasta S 2006 The economic effects of harmful algal blooms *Ecology of Harmful Algae* ed E Granéli and J T Turner (Dordrecht: Springer) pp 393–401
- Houlton B Z, Boyer E, Finzi A, Galloway J, Leach A, Liptzin D, Melillo J, Rosenstock T S, Sobota D and Townsend A R 2013 Intentional versus unintentional nitrogen use in the United States: trends, efficiency and implications *Biogeochemistry* 114 11–23
- Howarth R W 2008 Coastal nitrogen pollution: a review of sources and trends globally and regionally *Harmful Algae* 8 14-20
- Howarth R W and Paerl H 2008 Coastal marine eutrophication: control of both nitrogen and phosphorus is necessary *Proc. Natl Acad. Sci. USA* **105** E103
- Howarth R W, Swaney D P, Boyer E W, Marino R, Jaworski N and Goodale C 2006 The influence of climate on average nitrogen export from large watersheds in the Northeastern United States *Biogeochemistry* 79 163–86
- IFA(International Fertilizer Association) 2014 *IFA Database* (www. fertilizer.org/ifa/Home-Page/STATISTICS)
- Jeong H J, Yoo Y D, Kim T H, Kim J H, Kang N S and Yih W H 2004 Mixotrophy in the phototrophic harmful alga *Cochlodinium polykikoides* (dinophycean): prey species, the effects of prey concentration and grazing impact J. Eukaryot. *Microbiology* **51** 563–9
- Jewett E B, Lopez C B, Dortch Q, Etheridge S M and Baker L C 2008 Harmful Algal bloom management and response: assessment and Plan (Washington DC: Interagency Working Group on Harmful Algal Blooms, Hypoxia, and Human Health of the Joint Subcommittee on Ocean Science and Technology) pp 76
- Jickells T 2006 The role of air-sea exchange in the marine nitrogen cycle *Biogeosci. Disc.* **3** 183–210
- Johansson N and Granéli E 1999 Influence of different nutrient conditions on cell density, chemical composition and toxicity of *Prymnesium parvum* (Haptophyta) in semi-continuous cultures J. Exp. Mar. Biol. Ecol. 239 243–58
- Khakural B R and Alva A K 1995 Hydrolysis of urea in two sandy soils under citrus production as influenced by rate and depth of placement *Commun. Soil Sci. Plant Anal.* 26 2143–6
- Kosten S *et al* 2012 Warmer climates boost cyanobacterial dominance in shallow lakes *Glob. Change Biol.* **18** 118–26
- Kudela R M, Lane J Q and Cochlan W P 2008 The potential role of anthropogenically derived nitrogen in the growth of harmful algae in California, USA *Harmful Algae* 8 103–10
- Lee S J, Jang M H, Kim H S, Yoon B D and Oh H M 2000 Variation of microcystin content of *Microcystis aeruginosa* relative to medium N : P ratio and growth stage *J. Appl. Microbiology* **89** 323–9
- Leong S C Y, Murata A, Nagashima Y and Taguchi S 2004 Variability in toxicity of the dinoflagellate *Alexandrium tamarense* in response to different nitrogen sources and concentration *Toxicon* **43** 407–15
- Lewis W M Jr and Wurtsbaugh W A 2008 Control of lacustrine phytoplankton by nutrients: Erosion of the phosphorus paradigm *Int. Rev. Hydrobiol.* **93** 446–65

- Lewis W M Jr, Wurtsbaugh W A and Paerl H W 2011 Rationale for control of anthropogenic nitrogen and phosphorus to reduce eutrophication of inland waters *Environ. Sci. Technol.* **45** 10300–5
- Li J, Glibert P M, Zhou M, Lu S and Lu D 2009 Relationships between nitrogen and phosphorus forms and ratios and the development of dinoflagellate blooms in the East China Sea *Mar. Ecol. Prog. Ser.* 383 11–26
- Litke D W 1999 Review of phosphorus control measures in the United States and their effects on water quality US Geological Survey Water-Resources Investigations Report 99-4007 (Denver, CO: US Geological Survey)
- Lomas M W, Trice T M, Glibert P M, Bronk D A and McCarthy J J 2002 Temporal and spatial dynamics of urea uptake and regeneration rates and concentrations in Chesapeake Bay *Estuaries* 25 469–82
- McCarthy M J, James R T, Chen Y, East T L and Gardner W S 2009 Nutrient ratios and phytoplankton community structure in the large, shallow, eutrophic, subtropical lakes Okeechobee (FL, USA) and Taihu (China) *Limnology* **10** 215–27
- Monchamp M E, Pick F R, Beisner B E and Maranger R 2014 Variation in microcystin concentration and composition in relation to cyanobacterial community structure *PLoS One* **9** e85573
- Ning X, Lin C, Hao Q, Liu C, Le F and Shi J 2009 Long term changes in the ecosystem in the Northern South China Sea during 1976–2004 *Biogeoscice* 6 2227–43
- Nixon S W 1995 Coastal marine eutrophication: a definition, social causes, and future concerns *Ophelia* **41** 199–219
- Oh H-M, Lee S J, Jang M-H and Yoon B-D 2000 Microcystin production by *Microcystis aeruginosa* in a phosphorus-limited chemostat *Appl. Environ. Microbiology* **66** 176–9
- O'Neil J M, Davis T W, Burford M A and Gobler C J 2012 The rise of harmful cyanobacteria blooms: the potential roles of eutrophication and climate change *Harmful Algae* 14 313–34
- Orihel D M *et al* 2012 High microcystin concentrations occur only at low nitrogen-to-phosphorus ratios in nutrient-rich Canadian lakes *Can. J. Fish. Aquat. Sci.* **69** 1–6
- Otten T G, Xu H, Qin B, Zhu G and Paerl H W 2012 Spatiotemporal patterns and ecophysiology of toxigenic *Microcystis* blooms in Lake Taihu, China: implications for water quality management *Environ. Sci. Technol.* **46** 3480–8
- Paerl H W 2009 Controlling eutrophication along the freshwater–marine continuum: dual nutrient (N and P) reductions are essential *Estuaries Coasts* **32** 593–601
- Paerl H W and Otten T G 2013 Blooms bite the hand that feeds them Science 342 433–4
- Paerl H W and Scott J T 2010 Throwing fuel on the fire: synergistic effects of excessive nitrogen inputs and global warming on harmful algal blooms *Environ. Sci. Technol.* **44** 7756–8
- PBL Netherlands Environmental Assessment Agency 2012 Roads from RIO + 20. Pathways to Achieve Global Sustainability Goals by 2050 (The Hague: PBL Netherlands Environmental Assessment Agency) p 283 (www.pbl.nl/en/publications/2012/ roads-from-rio20)
- Peñuelas J, Sardans J, Rivas-Ubach A and Janssens I A 2012 The human-induced imbalance between C, N and P in Earth's life system *Glob. Change Biol.* 18 3–6
- Redfield A C 1934 On the proportions of organic derivatives in sea water and their relation to the composition of plankton *James Johnstone Memorial Volume* (Liverpool: University of Liverpool Press) pp 176–92
- Rhue R D and Harris W G 1999 Phosphorus sorption/desorption reactions in soil and sediments *Phosphorus Biogeochemistry in Subtropical Systems* ed K R Reddy, G A O'Connor and C L Schelske (Boca Raton, FL: Lewis Publishers) pp 187–206
- Rolland A, Bird D F and Giani A 2005 Seasonal changes in composition of the cyanobacterial community and the

occurrence of hepatotoxic blooms in the Eastern townships, Quebec, Canada J. Plankt. Res. 27 683–94

- Ryther J H and Dunstan W M 1971 Nitrogen, phosphorus and eutrophication in the marine environment *Science* **171** 1008–13
- Sattari S Z, Bouwman A F, Giller K E and van Ittersum M K 2012 Residual soil phosphorus as the missing piece in the global phosphorus crisis puzzle *Proc. Natl Acad. Sci. USA* 109 6348–54
- Sattari S Z, Van Ittersum M K, Giller K E, Zhang F and Bouwman A F 2014 Key role of China and its agriculture in global sustainable phosphorus management *Environ. Res. Lett.* 9 054003
- Schindler D W 1977 Evolution of phosphorus limitation in lakes Science 195 260–2
- Schindler D W, Hecky R E, Findlay D L, Stainton M P, Parker B R, Paterson M J, Beaty K G, Lyng M and Kasien S E M 2008 Eutrophication of lakes cannot be controlled by reducing nitrogen input: results of a 37-year whole-ecosystem experiment *Proc. Natl Acad. Sci. USA* 105 11254–8
- Scott J T and McCarthy M J 2010 Nitrogen fixation may not balance the nitrogen pool in lakes over timescales relevant to eutrophication management *Limnol. Oceanogr.* 55 1265–70
- Scott J T, McCarthy M J, Otten T G, Steffen M M, Baker B C, Grantz E M, Wilhelm S W and Paerl H W 2013 Comment: an alternative interpretation of the relationship between TN : TP and microcystins in Canadian lakes *Can. J. Fish. Aquat. Sci.* 70 1265–8
- Seitzinger S P 1988 Denitrification in freshwater and coastal marine ecosystems—ecological and geochemical significance *Limnol. Oceanogr.* **33** 702–24
- Shen Z-L and Liu Q 2009 Nutrients in the Changjiang River Environ. Monit. Assess. 153 27–44
- Shimizu Y, Watanabe N and Wrensford G 1993 Biosynthesis of brevetoxins and heterotrophic metabolism *Gymnodinium Breve Harmful Marine Algal Blooms* ed P Lassus, G Arzul, E Erard-Le-Denn, P Gentien and C Marcaillou (Paris: Lavoisier Publishing) pp 351–7
- Siuda W and Chrost R J 2006 Urea and ureolytic activity in lakes of different trophic status *Pol. J. Microbiology* **55** 211–25
- Sivonen K and Jones G J 1999 Cyanobacterial toxins Toxic Cyanobacteria in Water: A guide to their Public Health Consequences, Monitoring and Management ed I Chorus and J Bartram (London: Spon) pp 41–111
- Smil V 1999 Detonator of the population explosion Nature 400 415
- Smil V 2000 Phosphorous in the environment: natural flows and human interferences Ann. Rev. Energy. Environ. 25 25–53
- Smil V 2001 Enriching the Earth: Fritz Haber, Carl Bosch, and the Transformation of World Food (Cambridge, UK: MIT Press)
- Smith V H 1983 Low nitrogen to phosphorus ratios favor dominance by blue-green-algae in lake phytoplankton Science 221 669–71
- SOA (State Oceanographic Administration of China) 2010 Marine disaster communiqué of China (1989–2009) (www.mep.gov. cn/plan/zkgb.2010.8.13)
- Sobota D J, Compton J E and Harrison J A 2013 Reactive nitrogen inputs to US lands and waterways: how certain are we about sources and fluxes? *Frontiers Ecol. Environ.* **11** 82–90
- Solomon C M, Collier J L, Berg G M and Glibert P M 2010 Role of urea in microbes in aquatic systems: a biochemical and molecular review Aquat. Microb. Ecol. 59 67–88
- Sutton et al 2013 Our Nutrient World: The Challenge to Produce more Food and Energy with Less Pollution Global Overview of Nutrient Management. Centre for Ecology and Hydrology, Edinburgh on behalf of the Global Partnership on Nutrient Management and the International Nitrogen Initiative
- Syers J K, Johnston A A and Curtin D 2008 Efficiency of Soil and Fertilizer Phosphorus Use. Reconciling Changing Concepts of Soil Phophorous Behaviour with Agronomic Information (Rome: FAO) p 110

- Sylvan J B, Dortch Q, Nelson D M, Maier Brown A F, Morrison W and Ammerman J W 2006 Phosphorus limits phytoplankton growth on the Lousiana shelf during the period of hypoxia formation *Environ. Sci Technol.* **40** 7548–53
- Switzer T 2008 Urea loading from a spring storm-Knysna estuary, South Africa Harmful Algae **8** 66–9
- Ti C and Yan X 2013 Spatial and temporal variations of river nitrogen exports from major basins in China *Environ. Sci. Pollut. Res.* **20** 6509–20
- Van de Waal D B, Smith V H, Declerck S A, Stam E C M and Elser J J 2014 Stoichiometric regulation of phytoplankton toxins *Ecol. Lett.* 17 736–42
- Van de Waal D B, Verspagen J M H, Lurling M, Van Donk E, Visser P M and Huisman J 2009 The ecological stoichiometry of toxins produced by harmful cyanobacteria: an experimental test of the carbon-nutrient balance hypothesis *Ecol. Lett.* **12** 1326–35
- Van Drecht G, Bouwman A F, Harrison J and Knoop J M 2009 Global nitrogen and phosphate in urban waste water for the period 1970–2050 *Glob. Biogeochem. Cycles* 23 GB0A03
- Van Mooy B A S et al 2009 Phytoplankton in the ocean use nonphosphorus lipids in response to phosphorus scarcity Nature 458 69–72
- Vézie C, Rapala J, Vaitomaa J, Seitsonen J and Sivonen K 2002 Effect of nitrogen and phosphorus on growth of toxic and nontoxic *Microcystis* strains and on intracellular *Microcystin* concentrations *Microb. Ecol.* 43 443–54
- Wali P, Kumar V and Singh J P 2003 Effect of soil type, exchangeable sodium percentage, water content, and organic

amendments on urea hydrolysis in some tropical Indian soils Aust. J. Soil Res. 41 1171-6

- Wang B, Xie L and Sun X 2011 Water quality and marginal seas off China in the last two decades *Int. J. Oceanogr.* 2011 731828
- Wang H and Wang H 2009 Mitigation of lake eutrophication: loosen nitrogen control and focus on phosphorus abatement *Prog. Nat. Sci.* **19** 1445–51
- Wang S F, Tang D L, He F L, Fukuyo Y and Azanza R 2008 Occurrences of harmful algal blooms (HABs) associated with ocean environments in the South China Sea *Hydrobiologia* 596 79–93
- Xiao H-Y, Tang C-G, Xiao H-W, Liu X-Y and Liu C-Q 2010 Mosses indicating atmospheric nitrogen deposition in the Yangtze River drainage basin, China J. Geophys. Res. 115 D14301
- Xing G X and Zu Z L 2002 Regional nitrogen budgets for China and its major watersheds *Biogeochemistry* **57** 405–27
- Ye C, Xu Q, Kong H, Shen Z and Yan C 2007 Eutrophication conditions and ecological status in typical bays of Lake Taihu in China *Environ. Monit. Assess.* **135** 217–25
- Zhang W and Zhang F No Date The fertilizer Situation and Outlook in China (www.firt.org/sites/default/files/ Zhang_FertilizerMarketinChina_presentation.pdf) Accessed.22 June, 2014
- Zhou M J, Shen Z L and Yu R C 2008 Responses of a coastal phytoplankton community to increased nutrient input from the Changjiang (Yangtze) River *Cont. Shelf Res.* **28** 1483–9